

Mouse anti-Chicken IgY, clone 7C2, Biotin (Monoclonal)

Clone no. 7C2

MONOSAN

Product name	Mouse anti-Chicken IgY, clone 7C2, Biotin (Monoclonal)
Host	Mouse
Applications	ELISA,WB
Species reactivity	chicken, duck
Conjugate	Biotin
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	7C2
Size	1 ml
Concentration	50 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Chicken IgY, clone 7C2, Biotin (Monoclonal)

Clone no. 7C2

MONOSAN

Additional info

The monoclonal antibody 7C2 can be used during various purification steps of IgY. The yolk of eggs laid by immunized chickens has been recognized as an excellent source of polyclonal antibodies (pAb). Specific antibodies produced in chickens offer several important advantages over producing antibodies in other mammals. Because a single egg contains as much antibody as an average bleed from a rabbit, this simple, non-invasive approach presents an appealing alternative to conventional pAb production methods. Purification of chicken egg yolk immunoglobulin Y (IgY), the 150 kDa IgG homolog, does not require animal bleeding. In addition, the eggs from immunized chickens provide a continual, daily source of pAb, and this convenient approach offers greater compatibility with animal protection regulations. Due to the phylogenetic distance between birds and mammals, there is greater potential of producing a higher percentage of specific antibody against mammalian antigens when using chickens. Highly conserved mammalian proteins sometimes fail to illicit a humoral response in animals, such as rabbits, that are traditionally used for generating pAb. Non-specific binding and need for cross-species immunoabsorptions is eliminated since chicken IgY does not cross-react with mammalian IgG and does not bind bacterial or mammalian Fc receptors. There are well defined structural differences of IgY-type immunoglobulins and the IgG of mammals. That includes the molar mass of the heavy chains of the immunoglobulins. The IgY-type immunoglobulins are much less flexible than IgG. Also, the structures of the Fc part of the immunoglobulin isotypes IgY and IgG are different. The antibody 7C2 is cross reactive for duck IgY.

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES