

Mouse anti-PR3, clone PR3G-2 (Monoclonal)

Clone no. PRG-2

MONOSAN

Product name	Mouse anti-PR3, clone PR3G-2 (Monoclonal)
Host	Mouse
Applications	FC,FUNC,ELISA,WB
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	PRG-2
Size	1 ml
Concentration	100 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-PR3, clone PR3G-2 (Monoclonal)

Clone no. PRG-2

MONOSAN

Additional info

Monoclonal antibody PR3G-2 reacts with human proteinase 3 (PR3), a 30 kDa protein. PR3 is a major antigen recognized by autoantibodies directed against cytoplasmic proteins of neutrophilic granulocytes and monocytes (called anti-neutrophil cytoplasmic autoantibodies (ANCA)). ANCA are able to activate primed neutrophils to produce oxygen radicals and release lytic enzymes, including PR3. Proteinase 3 (PR3) was identified as the target antigen of ANCA in Wegener's granulomatosis (WG). ANCA directed against PR3 (PR3-ANCA) can interfere with the binding of PR3 to its physiological inhibitor alpha1-antitrypsin (alpha1-AT) and with the proteolytic activity of PR3. At the site of inflammation, PR3 can cleave the PR3-ANCA complex between these inhibiting ANCA and PR3 itself, leaving active PR3. Autoantibodies to PR3 are potent activators of the 5-lipoxygenase pathway in primed human neutrophils. Extracellular free arachidonic acid, as present at an inflammatory focus, synergizes with such autoantibodies to evoke full-blown lipid mediator generation, granule secretion and respiratory burst. Proteinase 3 (PR3) is a neutral serine proteinase, which is localized in the azurophilic granules of neutrophils and in granules of monocytes and can be detected in the membrane of secretory vesicles. PR3 degrades a number of extracellular matrix proteins such as elastin and inactivates human C1 inhibitor. Membrane-associated PR3 is also able to activate caspase-3 without triggering apoptosis of neutrophils, which is possibly a neutrophil survival mechanism. In addition, PR3 is involved in myeloid differentiation and is, therefore, also called myeloblastin. The monoclonal antibody PR3G-2 was produced by immunization of mice with a crude granule extract.

References

1. Geld van der; Y et al. Clin Exp Immunol 1999; 118: 487
2. Pederzoli, M et al: J Immunol 2005, 174: 6381
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES