Product datasheet

MON9846



MONOSAN

Mouse anti-IgA Secretory Component, clone SC-05 (Monoclonal)Clone no.SC-05

Mouse anti-IgA Secretory Component, clone SC-05 (Monoclonal)
Mouse
ELISA, FC, IHC-fr, IHC-P, IP, WB
human, rat
-
affinity purified human secretory component
lgG1-K
Monoclonal
SC-05
100 ug
100 ug/ml
-
PBS with 0.02% sodium azide
2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

www.monosan.com

Product datasheet

MON9846



Mouse anti-IgA Secretory Component, clone SC-05 (Monoclonal)Clone no.SC-05

MONOSAN

Additional info

SC-05 reacts with a reduction resistant epitope on 80 kDa human secretory component (both free and bound to SIgA). Secretory component is differentially expressed in epithelium, thus SC-05 can identify subpopulations of epithelial cells and epithelial differentiation. Secretory component negative cell lines are not stained with SC-05.

- 1. Kvale, D. et al, Int J Cancer 42(4): 638-641 (1988)
- 2 Bartek, J. et al, Histochem 91(3): 235-244 (1989)
- 3. Bartek, J. et al, Histochem J 22(10): 537-534 (1990)
- 4.
- 5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES