

Mouse anti-HGAL, clone MRQ-49 (Monoclonal)

Clone no. MRQ-49

MONOSAN

Product name	Mouse anti-HGAL, clone MRQ-49 (Monoclonal)
Host	Mouse
Applications	IHC-P (1:100-1:500)
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG2a-k
Clonality	Monoclonal
Clone number	MRQ-49
Size	1 ml
Concentration	n/a
Format	-
Storage buffer	Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-HGAL, clone MRQ-49 (Monoclonal)

Clone no. MRQ-49

MONOSAN

Additional info

Human germinal center associated lymphoma (HGAL) protein is specifically expressed in the cytoplasm of germinal center B-cells, but is absent in mantle and marginal zone B-cells and in the interfollicular and paracortical regions in normal tonsils and lymph nodes. Its high degree of specificity for germinal center B-cells makes anti-HGAL an ideal marker for the detection of germinal center-derived B-cell lymphomas. Anti-HGAL has the highest overall sensitivity of detecting follicular lymphoma (FL) and in detecting the interfollicular and diffuse components of FL compared with antibodies against bcl2, LMO2, CD10, and bcl6. The addition of anti-HGAL to the immunohistologic panel is beneficial in the work-up of nodal and extranodal B-cell lymphomas, and the efficacy of anti-HGAL in detecting the follicular, interfollicular, and diffuse components of FL is of particular value in the setting of variant immunoarchitectural patterns

References

1. Natkunam Y, et al. Blood. 2005; 105:3979–86
2. Natkunam Y, et al. Blood. 2007; 109:298-305
3. Younes SF, et al. Am J Surg Pathol. 2010; 34:1266-76
4. Higgins RA, et al. Arch Pathol Lab Med. 2008; 132:441-6
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES